



I. Epidemiological profile

Population	2014	%
High transmission (>1 case per 1000 population)	17 600 000	100
Low transmission (0-1 cases per 1000 population)	0	0
Malaria free (0 cases)	0	0
Total	17 600 000	

Parasites and vectors

Major plasmodium species: *P. falciparum* (100%), *P. vivax* (0%)
 Major anopheles species: *An. gambiae*, *An. funestus*, *An. arabiensis*
 Programme phase: Control

Reported confirmed cases: 5 428 655 Estimated cases, 2013: [4 700 000-10 000 000]
 Reported deaths: 5632 Estimated deaths, 2013: [12 000-32 000]

II. Intervention policies and strategies

Intervention	Policies/strategies	Yes/No	Adopted
ITN	ITNs/LLINs distributed free of charge	Yes	2007
	ITNs/LLINs distributed to all age groups	Yes	1998
IRS	IRS is recommended	Yes	2006
	DDT is authorized for IRS	No	-
Larval control	Use of larval control recommended	Yes	2012
IPT	IPT used to prevent malaria during pregnancy	Yes	2005
Diagnosis	Patients of all ages should receive diagnostic test	Yes	2009
	Malaria diagnosis is free of charge in the public sector	Yes	2009
Treatment	ACT is free for all ages in public sector	No	-
	Sale of oral artemisinin-based monotherapies	Never allowed	-
	Single dose of primaquine is used as gametocidal medicine for <i>P. falciparum</i>	No	-
	Primaquine is used for radical treatment of <i>P. vivax</i>	No	-
	G6PD test is a requirement before treatment with primaquine	No	-
	Directly observed treatment with primaquine is undertaken	No	-
	System for monitoring of adverse reactions to antimalarials exists	Yes	2009
Surveillance	ACD for case investigation (reactive)	No	-
	ACD of febrile cases at community level (pro-active)	No	-
	Mass screening is undertaken	No	-
	Uncomplicated <i>P. falciparum</i> cases routinely admitted	Yes	-
	Uncomplicated <i>P. vivax</i> cases routinely admitted	No	-

Antimalaria treatment policy

Medicine	Adopted
First-line treatment of unconfirmed malaria	AL; AS+AQ 2005
First-line treatment of <i>P. falciparum</i>	AL; AS+AQ 2005
Treatment failure of <i>P. falciparum</i>	QN -
Treatment of severe malaria	AS; QN -
Treatment of <i>P. vivax</i>	-
Dosage of primaquine for radical treatment of <i>P. vivax</i>	-
Type of RDT used	<i>P. f</i> only.



Therapeutic efficacy tests (clinical and parasitological failure, %)

Medicine	Year	Min	Median	Max	Follow-up	No. of studies	Species
AL	2005-2012	0	6.15	12.5	28 days	9	<i>P. falciparum</i>
AS+AQ	2006-2012	0	5.05	21.5	28 days	6	<i>P. falciparum</i>





Insecticide susceptibility bioassays (reported resistance to at least one insecticide for any vector at any locality)

Year	Pyrethroid	DDT	Carbamate	Organophosphate	Species/complex tested
2010-2014	Yes	Yes	Yes	Yes	<i>An. arabiensis</i> , <i>An. coluzzii</i> , <i>An. gambiae</i> s.l.



III. Financing



IV. Coverage



V. Impact



Impact: Insufficiently consistent data to assess trends